

OBSERVATIONS REGARDING COMPACTNESS IN THE $\bar{\partial}$ -NEUMANN PROBLEM

MEHMET ÇELİK & EMIL J. STRAUBE

1. INTRODUCTION

Compactness of the $\bar{\partial}$ -Neumann operator plays an important role in several contexts. The condition was initially introduced by Kohn and Nirenberg [22] as a sufficient condition for global regularity. Work of Catlin [6] and Sibony [31] showed that this does indeed provide a viable route to global regularity; that is, the compactness condition can be verified on large classes of domains. We refer the reader to [5, 9, 17, 33, 34] for background on the $\mathcal{L}^2$ -Sobolev theory of the $\bar{\partial}$ -Neumann problem in general and on compactness in particular.

Subellipticity of the $\bar{\partial}$ -Neumann operator is independent of the metric (a fact usually attributed to Sweeney ([35]), although that reference deals with the coercive case), while continuity in Sobolev spaces is not. The latter is a consequence of Kohn's results concerning estimates for the $\bar{\partial}$ -Neumann problem with weights ([20]) and Barrett's results on failure of Sobolev estimates on the worm domains ([2]). Compactness is intermediate between subellipticity and continuity in Sobolev spaces. Consequently, it is of interest to know that compactness is also independent of the metric, for metrics subject only to the condition that they be smooth on the closure of the domain (so that the induced norms on the L^2 -spaces of forms are equivalent). In particular, the metrics at higher form levels are not required to be induced by the metric on $(0, 1)$ -forms. We also obtain a new proof of the independence of subellipticity of the $\bar{\partial}$ -Neumann operator from the metric (for the same class of metrics).

We define the notion of a compactness multiplier in obvious analogy to that of a subelliptic multiplier ([21, 13]). It is easily seen that the continuous multipliers form an ideal of the form $\{f \in C(\bar{\Omega}) \mid f(z) = 0, z \in A\}$, where the common zero set A is a compact subset of the boundary. This common zero set may thus be viewed as the obstruction to compactness: the $\bar{\partial}$ -Neumann operator is compact if and only if A is the empty set. Of course, this purely abstractly defined obstruction is of use only to the extent that it can actually be determined if the domain is given. We do this for two classes of domains where compactness is understood, i.e. convex domains in $\mathbb{C}^n$ and complete Hartogs domains in $\mathbb{C}^2$. For a convex domain, and for $(0, 1)$ -forms, this set is the closure of the union of all the analytic discs in the boundary. For a Hartogs domain in $\mathbb{C}^2$, on the other hand, the closure of the union of all the analytic discs

2000 *Mathematics Subject Classification*. Primary 32W05; Secondary 35N15.

Work supported in part by NSF grants DMS-0500842 and DMS-0758534.

in the boundary can be strictly contained in A . This is a reflection of the fact that even in $\mathbb{C}^2$, there can be obstructions to compactness more subtle than analytic discs in the boundary ([25], [17], Theorem 4.2). What matters are fine interior points of the projection (on the base) of the weakly pseudoconvex points, not just Euclidean interior points of this projection (which correspond to analytic discs in the boundary, cf. [34], Lemma 3.18). Accordingly, A equals the Euclidean closure of the inverse image of these fine interior points (under a mild technical condition).

Much of this material comes from the first author's Ph.D. dissertation ([8]) written at Texas A&M University under the supervision of the second author.

2. VARIATION OF THE METRIC

A theorem by W. J. Sweeney in [35] shows that coercive estimates are independent of the metric on the tangent bundle, and the fact that the same is true for subelliptic estimates is also usually attributed to him. In view of Kohn's results concerning Sobolev estimates for the $\bar{\partial}$ -Neumann operator associated to suitably weighted metrics ([20]), and Barrett's results on failure of Sobolev estimates on the worm domains ([2]), this invariance does not hold for Sobolev estimates. As compactness is a regularity property that lies between subellipticity and continuity in Sobolev spaces, it is natural to ask how it behaves when the metric is changed, and we note that it is independent of the metric. We then give a new proof that subellipticity of the $\bar{\partial}$ -Neumann operator is independent of the metric.

Let Ω be a bounded pseudoconvex domain in $\mathbb{C}^n$. For $q = 1, \dots, n$, denote by $G^q(z) = G_{I,J}^q(z)$ a smooth function on $\bar{\Omega}$ with values in the strictly positive definite Hermitian $(n!/(n-q)!q!)$ by $(n!/(n-q)!q!)$ matrices; I and J are strictly increasing q -tuples. We denote by $L_{(0,q)}^2(\Omega, G^q)$ the square-integrable $(0, q)$ -forms, but with the standard inner product replaced by the one given pointwise by $G_{I,J}^q(z)$. Let $u = \sum'_{|J|=q} u_J d\bar{z}_J$, $v = \sum'_{|K|=q} v_K d\bar{z}_K$. Here, the prime indicates as usual summation over increasing q -tuples, and $d\bar{z}_J = d\bar{z}_{j_1} \wedge \dots \wedge d\bar{z}_{j_q}$. Then

$$(1) \quad (u, v)_{G^q} := \int_{\Omega} \langle u, v \rangle_{G^q} = \sum'_{|J|=|K|=q} \int_{\Omega} G_{JK}^q(z) u_J(z) \overline{v_K(z)} dV,$$

and

$$(2) \quad \|u\|_{G^q}^2 := \int_{\Omega} \langle u, u \rangle_{G^q} = \sum'_{|J|=|K|=q} \int_{\Omega} G_{JK}^q u_J(z) \overline{u_K(z)} dV.$$

Note that since $G_{I,J}^q$ is smooth on the closure of Ω and takes only positive definite matrices as values, the *set* of square-integrable forms does not change, but the inner product does (to an equivalent one). With this inner product, $L_{(0,q)}^2(\Omega, G^q)$ is a Hilbert space. It will be convenient to refer to the inner product in (1) and the norm in (2) as 'weighted'.

Remark 1: The metrics G^q for different q 's are not assumed related. In particular, it is not assumed that G^q is induced by G^1 .

We briefly recall the setup for the $\bar{\partial}$ -complex and the $\bar{\partial}$ -Neumann problem in the case of a general metric. First,

$$(3) \quad \bar{\partial}u = \bar{\partial} \left(\sum'_{|J|=q} u_J d\bar{z}_J \right) = \sum'_{|J|=q} \bar{\partial}u_J \wedge d\bar{z}_J$$

$$(4) \quad = \sum_{j=1}^n \sum'_{|J|=q} \frac{\partial u_J}{\partial \bar{z}_j} d\bar{z}_j \wedge d\bar{z}_J,$$

where the domain consists of those forms where the right hand side, when computed as distributions, is in $L^2_{(0,q+1)}(\Omega, G^{q+1})$. Thus $\bar{\partial}$ is closed and densely defined, at each level q , and so has a Hilbert space adjoint $\bar{\partial}_G^*$. For example, when $q = 0$, the usual computation for this adjoint gives

$$(5) \quad (\bar{\partial})_G^* u = - \sum_{j,k=1}^n (G^0)^{-1} \frac{\partial (G^1_{jk} u_j(z))}{\partial z_k}, \quad u \in \text{Dom} \left((\bar{\partial})_G^* \right);$$

the boundary condition is

$$(6) \quad \sum_{j,k=1}^n G^1_{jk} u_j \frac{\partial \rho}{\partial z_k} = 0 \quad \text{for } z \in b\Omega.$$

Note that if G^1 is the Euclidean metric, so that $G^1_{jk} = \delta_{jk}$ (where δ_{jk} is the Kronecker- δ), we obtain the familiar boundary condition $\sum_{k=1}^n u_k \frac{\partial \rho}{\partial z_k} = 0$ on $b\Omega$.

For $u, v \in \text{Dom}(\bar{\partial}_q) \cap \text{Dom} \left((\bar{\partial}_{q-1})_G^* \right)$, the Dirichlet form $Q_{G,q}(u, v)$ is defined as

$$(7) \quad Q_{G,q}(u, v) := (\bar{\partial}u, \bar{\partial}v)_{G^{q+1}} + \left((\bar{\partial}_{q-1})_G^* u, (\bar{\partial}_{q-1})_G^* v \right)_{G^{q-1}}.$$

$\text{Dom}(\bar{\partial}_q) \cap \text{Dom} \left((\bar{\partial}_{q-1})_G^* \right)$ is complete with respect to $\|u\|_G^2 := Q_G(u, u) + \|u\|_G^2$. Thus there is a unique non-negative selfadjoint operator $\square_q^G$ associated to $Q_{G,q}$ via

$$(8) \quad Q_{G,q}(u, v) = (\square_q^G u, v)_{G^q}, \quad u \in \text{Dom}(\square_q^G),$$

where $\text{Dom}(\square_q^G)$ consists of those u in $\text{Dom}(\bar{\partial}_q) \cap \text{Dom} \left((\bar{\partial}_{q-1})_G^* \right)$ with $\bar{\partial}u \in \text{Dom} \left((\bar{\partial}_q)_G^* \right)$ and $(\bar{\partial}_{q-1})_G^* u \in \text{Dom}(\bar{\partial}_{q-1})$. (See [30], Theorem VIII.15, or [14] Theorem 4.4.2: if Q is a closed symmetric quadratic form then Q is the quadratic form of a unique self adjoint operator as in (8)).

Because G^{q+1} induces a norm on $L^2_{(0,q+1)}(\Omega, G^{q+1})$ that is equivalent to the Euclidean one, the domain and range of $\bar{\partial}$ are unchanged from the Euclidean setting. In particular, the range is still equal to the kernel of $\bar{\partial}$ acting on $(q+1)$ -forms. The range of $(\bar{\partial}_q)_G^*$ is then also closed (because the range of $\bar{\partial}_q$ is, see for example, Lemma 4.1.1 in [9]). However, the range of $(\bar{\partial}_q)_G^*$ is also dense in $\ker(\bar{\partial}_q)^{\perp_{G^q}}$, and so

$\text{Im} \left((\bar{\partial}_q)_G^* \right) = \ker(\bar{\partial}_q)^{\perp_{G^q}}$. It follows that

$$(9) \quad L_{(0,q)}^2(\Omega, G^q) = \underbrace{\ker(\bar{\partial}_q)}_{\text{Im}(\bar{\partial}_{q-1})} \oplus \underbrace{\text{Im} \left((\bar{\partial}_q)_G^* \right)}_{\ker((\bar{\partial}_{q-1})_G^*)}$$

(since also $\text{Im}(\bar{\partial}_{q-1})^{\perp_{G^q}} = \ker((\bar{\partial}_{q-1})_G^*)$). Therefore, $\ker(\square_q^G) = \ker(\bar{\partial}_q) \cap \ker \left((\bar{\partial}_{q-1})_G^* \right) = \{0\}$ (the first equality is from (8)). Then, Theorem 1.1.2 in [19] implies

$$(10) \quad \|u\|_{G^q}^2 \lesssim \|\bar{\partial}_q u\|_{G^{q+1}}^2 + \|(\bar{\partial}_{q-1})_G^* u\|_{G^{q-1}}^2$$

for $u \in \text{Dom}(\bar{\partial}) \cap \text{Dom} \left((\bar{\partial}_{q-1})_G^* \right)$. (10) is the crucial estimate for the L^2 -theory, and general Hilbert space arguments now give as usual that $\square_q^G$ has a bounded inverse. In fact, if $u, v \in \text{Dom}(\bar{\partial}_q) \cap \text{Dom} \left((\bar{\partial}_{q-1})_G^* \right)$, then

$$(11) \quad |(u, v)_{G^q}| \leq \|u\|_{G^q} \|v\|_{G^q} \lesssim \|u\|_{G^q} \left(\|\bar{\partial}_q v\|_{G^{q+1}}^2 + \|(\bar{\partial}_{q-1})_G^* v\|_{G^{q-1}}^2 \right)^{\frac{1}{2}}.$$

That is, the functional on the left hand side of (11) is a continuous (conjugate linear) functional in v (for u fixed) in the norm induced by Q_{G^q} on $\text{Dom}(\bar{\partial}_q) \cap \text{Dom} \left((\bar{\partial}_{q-1})_G^* \right)$. Thus, it is given by an inner product

$$(12) \quad (u, v)_{G^q} = Q_{G,q}(N_q^G u, v).$$

N_q^G is the weighted $\bar{\partial}$ -Neumann operator. By definition, N_q^G maps $L_{(0,q)}^2(\Omega, G^q)$ continuously into $\text{Dom}(\bar{\partial}_q) \cap \text{Dom} \left((\bar{\partial}_{q-1})_G^* \right)$, a fortiori (by (10)) into $L_{(0,q)}^2(\Omega, G^q)$. It is immediate from (8) and (12) that N_q^G inverts $\square_q^G$. Denote by N_q the $\bar{\partial}$ -Neumann operator in the Euclidean metric.

Theorem 1. *Let Ω be a bounded pseudoconvex domain in $\mathbb{C}^n$, $1 \leq q \leq n$. Then N_q^G is compact if and only if N_q is compact.*

Proof. Both N_q and N_q^G can be expressed in terms of the canonical solution operators to $\bar{\partial}$:

$$(13) \quad N_q = \left((\bar{\partial}_{q-1})^* N_q \right)^* \left((\bar{\partial}_{q-1})^* N_q \right) + \left((\bar{\partial}_q)^* N_{q+1} \right) \left((\bar{\partial}_q)^* N_{q+1} \right)^*,$$

and

$$(14) \quad N_q^G = \left((\bar{\partial}_{q-1})_G^* N_q^G \right)_G^* \left((\bar{\partial}_{q-1})_G^* N_q^G \right) + \left((\bar{\partial}_q)_G^* N_{q+1}^G \right) \left((\bar{\partial}_q)_G^* N_{q+1}^G \right)_G^*.$$

For N_q , this is a well known fact, see [15], p.55, [28]; for N_q^G the proof is the same. Denote by P_q^G the orthogonal projection from $L_{0,q}^2(\Omega, G^q)$ onto $\ker(\bar{\partial}_q)$. Since $(\bar{\partial}_{q-1})_G^* N_q^G$ annihilates $\ker(\bar{\partial}_q)^{\perp_{G^q}}$, we have

$$(15) \quad (\bar{\partial}_{q-1})_G^* N_q^G = (\bar{\partial}_{q-1})_G^* N_q^G P_q^G.$$

Now, if f is a $\bar{\partial}$ -closed $(0, q)$ -form, then $\bar{\partial}^* N_q f$ and $(\bar{\partial}_{q-1})_G^* N_q^G f$ are both solutions of the equation $\bar{\partial} u = f$; orthogonal in the respective inner products to $\ker(\bar{\partial}_{q-1})$. Therefore the previous formula implies

$$(16) \quad (\bar{\partial}_{q-1})_G^* N_q^G = (I - P_{q-1}^G) \bar{\partial}^* N_q P_q^G ,$$

and (with $q+1$ in place of q)

$$(17) \quad (\bar{\partial}_q)_G^* N_{q+1}^G = (I - P_q^G) \bar{\partial}^* N_{q+1} P_{q+1}^G .$$

Analogously,

$$(18) \quad \bar{\partial}^* N_q = (I - P_{q-1}) (\bar{\partial}_{q-1})_G^* N_q^G P_q ,$$

and

$$(19) \quad \bar{\partial}^* N_{q+1} = (I - P_q) (\bar{\partial}_q)_G^* N_{q+1}^G P_{q+1} .$$

By using the fact that $A^*A + BB^*$ is compact if and only if A and B are compact, the above identities, and the fact that composition with bounded operators (projections in our case) preserves compactness, we obtain the theorem. $\square$

For the rest of this section, we assume that Ω is also (C^∞) smooth. A subelliptic estimate of order $\varepsilon > 0$ is said to hold for the $\bar{\partial}$ -Neumann problem, if

$$(20) \quad \|u\|_{G^q, \varepsilon}^2 \leq C \left(\|\bar{\partial} u\|_{G^{q+1}}^2 + \|\bar{\partial}^* u\|_{G^{q-1}}^2 \right), \quad u \in C_{(0,q)}^\infty(\bar{\Omega}) \cap \text{Dom} \left((\bar{\partial}_{q-1})_G^* \right),$$

where the norm on the left hand side is the L^2 -Sobolev norm of order ε . We remark that integer Sobolev norms are defined as weighted L^2 -norms of derivatives of forms (acting coefficientwise), and the noninteger ones are then obtained by interpolation. A subelliptic estimate holds if and only if N_q^G maps $L_{(0,q)}^2(\Omega)$ continuously to $W_{(0,q)}^{2\varepsilon}(\Omega)$. The proof in the weighted case is the same as in the Euclidean case.

Theorem 2. *Let Ω be a smooth bounded pseudoconvex domain in $\mathbb{C}^n$, $1 \leq q \leq n$, $\varepsilon > 0$. Then N_q^G is subelliptic of order 2ε if and only if N_q is subelliptic of order 2ε .*

Proof. By the Riesz representation theorem, there is an isomorphism $T_q^G : L_{(0,q)}^2(\Omega) \rightarrow L_{(0,q)}^2(\Omega, G^q)$ such that

$$(21) \quad (u, v) = (T_q^G u, v)_{G^q} .$$

T_q^G can also be computed directly from (1); of importance for us is the fact that the coefficients of $T_q^G u$ are linear combinations, whose coefficients are functions in $C^\infty(\bar{\Omega})$, of the coefficients of u . A direct computation shows that $u \in \text{Dom}(\bar{\partial}_{q-1}^*)$ if and only if $T_q^G u \in \text{Dom} \left((\bar{\partial}_{q-1})_{G^q}^* \right)$, and that

$$(22) \quad (\bar{\partial}_{q-1})_{G^q}^* T_q^G u = T_{q-1}^G (\bar{\partial}_{q-1})^* u .$$

First assume that there is a subelliptic estimate of order $\varepsilon > 0$ in the weighted norm. Let $u \in \text{Dom}(\bar{\partial}_q) \cap \text{Dom}(\bar{\partial}_q^*)$. Then $T_q^G u \in \text{Dom}\left((\bar{\partial}_{q-1})_{G_q}^*\right)$, and we have

$$\begin{aligned}
 (23) \quad \|u\|_\varepsilon^2 &\lesssim \|T_q^G u\|_{\varepsilon, G^q}^2 + \|u\|^2 \\
 &\lesssim \|\bar{\partial}_q T_q^G u\|_{G^{q+1}}^2 + \|(\bar{\partial}_{q-1})_G^* T_q^G u\|_{G^{q-1}}^2 + \|u\|^2 \\
 &\lesssim \sum'_{|K|=q} \sum_{j=1}^n \left\| \frac{\partial u_K}{\partial \bar{z}_j} \right\|^2 + \|u\|^2 + \|T_{q-1}^G \bar{\partial}_{q-1}^* u\|^2 \\
 &\lesssim \sum'_{|K|=q} \sum_{j=1}^n \left\| \frac{\partial u_K}{\partial \bar{z}_j} \right\|^2 + \|u\|^2 + \|\bar{\partial}_{q-1}^* u\|^2 \lesssim \|\bar{\partial}_q u\|^2 + \|\bar{\partial}_{q-1}^* u\|^2.
 \end{aligned}$$

The third inequality results from the form of $T_q^G u$ pointed out above, and the last inequality is from the Kohn-Morrey formula (see e.g. [9]). (23) is the desired subelliptic estimate for the unweighted metric.

The above argument relies (among other things) on the Kohn-Morrey inequality. Instead of attempting to derive a (complicated) version for the case of general metrics G^q , $1 \leq q \leq n$, we give a different argument for the proof of the reverse direction (this argument could also be used for the first direction). When $q = 1$, this argument will involve the $\bar{\partial}$ -Neumann operators N_0 and N_0^G ; a reference is [9] Theorem 4.4.3.

Let $u \in \text{Dom}(\bar{\partial}_q) \cap \text{Dom}\left((\bar{\partial}_{q-1})_{G_1}^*\right) \subset L^2_{(0,q)}(\Omega, G^q)$. Then

$$(24) \quad u = (\bar{\partial}_q)_G^* N_{q+1}^G (\bar{\partial}_q u) + \bar{\partial}_{q-1} N_{q-1}^G \left((\bar{\partial}_{q-1})_G^* u \right).$$

Because $(I - P_1^G)$ projects onto the range of $(\bar{\partial}_q)_{G_{q+1}}^*$, we have

$$(25) \quad (\bar{\partial}_q)_{G_{q+1}}^* N_{q+1}^G (\bar{\partial}_q u) = (I - P_q^G) \bar{\partial}_q^* N_{q+1} (\bar{\partial}_q u).$$

For the second term on the right of (24), we use that $\bar{\partial}_{q-1} N_{q-1}^G = N_q^G \bar{\partial}_{q-1}$ and then (25) with $q - 1$ in place of q to obtain

$$\begin{aligned}
 (26) \quad \bar{\partial}_{q-1} N_{q-1}^G &= N_q^G \bar{\partial}_{q-1} = \left((\bar{\partial}_{q-1})_G^* N_q^G \right)_G^* \\
 &= \left((I - P_{q-1}^G) \bar{\partial}_{q-1}^* N_q P_q^G \right)_G^* = P_q^G \left(\bar{\partial}_{q-1}^* N_q \right)_G^* (I - P_{q-1}^G).
 \end{aligned}$$

Thus, by using (25) and (26), we write (24) as

$$(27) \quad u = (I - P_q^G) \bar{\partial}_q^* N_{q+1} (\bar{\partial}_q u) + P_q^G \left(\bar{\partial}_{q-1}^* N_q \right)_G^* \left((\bar{\partial}_{q-1})_{G_q}^* u \right).$$

We have used that $(I - P_{q-1}^G) (\bar{\partial}_{q-1})_G^* u = (\bar{\partial}_{q-1})_G^* u$.

A subelliptic estimate associated with the metric G will follow if we can show the following two mapping properties (as continuous maps):

- (i) $P_q^G : W_{(0,q)}^\varepsilon(\Omega, G^q) \longrightarrow W_{(0,q)}^\varepsilon(\Omega, G^q)$,
- (ii) $\left(\bar{\partial}_{q-1}^* N_q \right)_G^* : L^2_{(0,q)}(\Omega, G^q) \longrightarrow W_{(0,q-1)}^\varepsilon(\Omega, G^q)$.

For (i), note that N_q is compact (since it is subelliptic); hence so is N_q^G , by Theorem 1. This implies that $P_q^G = \bar{\partial}\bar{\partial}_G^* N_q^G$ preserves the Sobolev spaces: the proof is analogous to the unweighted case ([9], Theorem 6.2.2). Note that the form $Q_{G,q}$ is also covered by the results in [22].

As for (ii), note that $W_{(0,q)}^\varepsilon(\Omega, G) = W_{(0,q)}^\varepsilon(\Omega)$ and $\left(W_{(0,q)}^\varepsilon(\Omega)\right)^* = W_{(0,q)}^{-\varepsilon}(\Omega)$, since $0 \leq \varepsilon \leq 1/2$; the spaces $W^\varepsilon(\Omega)$ and $W_0^\varepsilon(\Omega)$ coincide for $0 \leq \varepsilon \leq 1/2$ (with equivalent norms), see Theorem 11.1 in [24]. This duality similarly holds for the weighted spaces, with the weighted L^2 pairing. Thus, the statement in (ii) is equivalent to having an extension of $\bar{\partial}^* N_q$ as a continuous map

$$(28) \quad \bar{\partial}^* N_q : W_{(0,q)}^{-\varepsilon}(\Omega) \longrightarrow L_{(0,q-1)}^2(\Omega) .$$

This is a well know consequence of subellipticity of N_q , as follows. $L_{(0,q)}^2(\Omega)$ is dense in $W_{(0,q)}^{-\varepsilon}(\Omega)$, so to prove (28) we let $u \in L_{(0,q)}^2(\Omega)$ and estimate

$$(29) \quad \begin{aligned} \|\bar{\partial}^* N_q u\|^2 + \|\bar{\partial} N_q u\|^2 &= \left(\bar{\partial}\bar{\partial}^* N_q u, N_q u\right) + \left(\bar{\partial}^* \bar{\partial} N_q u, N_q u\right) \\ &= (u, N_q u) \lesssim \|u\|_{-\varepsilon} \|N_q u\|_\varepsilon \leq (\text{s.c.}) \|N_q u\|_\varepsilon^2 + (\text{l.c.}) \|u\|_{-\varepsilon}^2 \\ &\lesssim (\text{s.c.}) \left(\|\bar{\partial}^* N_q u\|^2 + \|\bar{\partial} N_q u\|^2\right) + (\text{l.c.}) \|u\|_{-\varepsilon}^2 . \end{aligned}$$

The last inequality comes from the subelliptic estimate associated with the Euclidean metric. The first term on the right in the last inequality can be absorbed into the left side of (29) to obtain

$$(30) \quad \|\bar{\partial}^* N_q u\|^2 + \|\bar{\partial} N_q u\|^2 \lesssim \|u\|_{-\varepsilon}^2 .$$

This was for $u \in L_{(0,q)}^2(\Omega)$. By density both $\bar{\partial}^* N_q$ and $\bar{\partial} N_q$ extend to continuous operators from $W_{(0,q)}^{-\varepsilon}(\Omega)$ to $L_{(0,q-1)}^2(\Omega)$ and $L_{(0,q+1)}^2(\Omega)$, respectively. In particular, (28) holds. This completes the proof of Theorem 2. $\square$

3. OBSTRUCTIONS TO COMPACTNESS

Let Ω be a bounded pseudoconvex domain in $\mathbf{C}^n$. Recall that N_q is compact if and only if there is a so called compactness estimate: for every $\varepsilon > 0$ there is a constant $C_\varepsilon > 0$ such that the estimate

$$(31) \quad \|u\|^2 \leq \varepsilon \left(\|\bar{\partial} u\|^2 + \|\bar{\partial}^* u\|^2\right) + C_\varepsilon \|u\|_{-1}^2$$

is valid for all $u \in \text{Dom}(\bar{\partial}) \cap \text{Dom}(\bar{\partial}^*) \subset L_{(0,q)}^2(\Omega)$ ([17], Lemma 1.1, [34], Proposition 3.2).

A function $f \in C(\bar{\Omega})$ is called a compactness multiplier on Ω if for every $\varepsilon > 0$ there is a constant $C_{\varepsilon,f} > 0$ such that the estimate

$$(32) \quad \|fu\|^2 \leq \varepsilon (\|\bar{\partial} u\|^2 + \|\bar{\partial}^* u\|^2) + C_{\varepsilon,f} \|u\|_{-1}^2$$

is valid for all $u \in \text{Dom}(\bar{\partial}) \cap \text{Dom}(\bar{\partial}^*) \subset L_{(0,q)}^2(\Omega)$. Note that $f \in C(\bar{\Omega})$ is a compactness multiplier if and only if the multiplication operator $M_f : u \rightarrow fu$ from

$\text{Dom}(\bar{\partial}) \cap \text{Dom}(\bar{\partial}^*)$, equipped with the graph norm, to $L^2_{(0,q)}(\Omega)$ is compact. Namely, in terms of the graph norm $\|u\|_{\text{graph}}$, estimate (32) says that $\|M_f u\|^2 \leq \varepsilon \|u\|_{\text{graph}}^2 + C_{\varepsilon,f} \|u\|_{-1}^2$. Because $L^2(\Omega)$ embeds compactly into $W^{-1}(\Omega)$, having this inequality for all $\varepsilon > 0$ characterizes compactness of the operator $M_f : \text{Dom}(\bar{\partial}) \cap \text{Dom}(\bar{\partial}^*) \rightarrow L^2_{(0,q)}(\Omega)$, see for example [12], Proposition V.2.3, [26], Lemma 2.1, [34], Lemma 3.3.

The basic properties of compactness multipliers are rather more elementary than the corresponding facts for subelliptic multipliers ([11], [13]). Let J^q be the set of the compactness multipliers defined as above, associated with $(0, q)$ forms, $1 \leq q \leq n$. Denote by A_q the common zero set of the elements of J^q . A_q is compact, and by interior elliptic regularity of the complex $\bar{\partial} \oplus \bar{\partial}^*$, $A_q \subseteq b\Omega$. More precisely, any $\varphi \in C_0^\infty(\Omega)$ is a compactness multiplier: if $u \in \text{Dom}(\bar{\partial}) \cap \text{Dom}(\bar{\partial}^*)$, then φu has components in $W_0^1(\Omega)$. The latter space embeds compactly into $L^2(\Omega)$, so that M_φ is indeed compact.

Proposition 1. *Let Ω be a bounded pseudoconvex domain in $\mathbb{C}^n$. The set of compactness multipliers J^q is a closed ideal in $C(\bar{\Omega})$, and so equals $\{f \in C(\bar{\Omega}) \mid f \equiv 0 \text{ on } A_q\}$.*

Proof. It is easy to see that hg is a compactness multiplier whenever g is; $\|(hg)u\|^2 \leq (\sup_{z \in \bar{\Omega}} |h(z)|) \|gu\|^2$. Thus, J^q is closed under multiplication by elements of $C(\bar{\Omega})$. The sum of two compactness multipliers is a compactness multiplier: $\|(g+f)u\|^2 \leq 2(\|gu\|^2 + \|fu\|^2)$. So J^q is an ideal of $C(\bar{\Omega})$.

To see that J^q is closed under the *sup*-norm, observe that the operator norm of M_f (as an operator from $\text{Dom}(\bar{\partial}) \cap \text{Dom}(\bar{\partial}^*) \rightarrow L^2_{(0,q)}(\Omega)$) is dominated by $\sup_{z \in \bar{\Omega}} |f(z)|$. Indeed, we have

$$(33) \quad \|M_f u\|^2 \leq \left(\sup_{z \in \bar{\Omega}} |f(z)| \right)^2 \|u\|^2 \leq \frac{D^2 e}{q} \left(\sup_{z \in \bar{\Omega}} |f(z)| \right)^2 \left(\|\bar{\partial} u\|^2 + \|\bar{\partial}^* u\|^2 \right),$$

where D is the diameter of Ω . The second inequality is the fundamental L^2 estimate for the $\bar{\partial}$ -complex dating back to Hörmander ([19], [9], [34]). Therefore, if $f \in C(\bar{\Omega})$ is a uniform limit of a sequence of compactness multipliers $\{f_n\}_{n=1}^\infty$, then the corresponding compact multiplication operators M_{f_n} converge in operator norm to M_f . Consequently, M_f is compact as well, and f is a compactness multiplier.

Finally, any closed ideal in $C(\bar{\Omega})$ is the full ideal generated by the zero set. For this elementary fact, see for example [23], Theorem 2.1. In our situation, this fact is also easily established directly. $\square$

Remark 2: Any function in $C(\bar{\Omega})$ that vanishes on the boundary is thus a compactness multiplier.

Remark 3: When Ω is a smooth domain, the set A_q is a subset of the set of boundary points of infinite type. This is immediate because a subelliptic pseudolocal estimate holds near a point of finite type ([7]).

Remark 4: If the set A_q is not empty, it cannot be ‘too small’. In particular, it cannot satisfy property (P_q) (see [6] for (P_1) , [17] and [34] for (P_q) , $q \geq 1$). The proof of this fact is analogous to the proof that compactness is a local property, see

Lemma 1.2 in [17], Proposition 3.4 in [34]; essentially the same argument also occurs in the first part of the proof of Theorem 3 below. One shows indirectly that A_q satisfying (P_q) implies a compactness estimate by writing a form u as $u_1 + u_2$, with u_1 supported near A_q , and u_2 supported away from A_q . Then u_1 is estimated by using the estimate $\sum_K' \sum_{j,k} \int_{\Omega} (\partial^2 \lambda / \partial z_j \partial \bar{z}_k) u_{j,K} \overline{u_{k,K}} \leq e(\|\bar{\partial} u\|^2 + \|\bar{\partial}^* u\|^2)$ ([5], p.83, [34], Corollary 1.12) in the usual way. u_2 is estimated via $u_2 = \varphi u_2$, where φ is supported away from A_q and so is a compactness multiplier. Details of this argument are in [8] and in the first part of the proof of Theorem 3 below. In particular, A_1 cannot have two dimensional Hausdorff measure zero, as such sets satisfy (P_1) ([3, 31]), nor can it be contained in a subvariety of the boundary of holomorphic dimension zero ([31], Proposition 12).

We do not attempt to develop a serious theory of compactness multipliers here; in particular, we ignore questions relating to the algorithmic point of view in [21, 13, 27]. Instead, we determine the sets A_q for two classes of domains.

Denote by $\{f_{\alpha}(\mathbb{D}^q)\}_{\alpha \in \Lambda}$ the family of q -dimensional analytic polydiscs in the boundary of Ω . That is, f_{α} is holomorphic on the q -dimensional unit polydisc $\mathbb{D}^q$ and continuous on its closure, and it maps into $b\Omega$. It was shown by Fu and Straube ([16]) that on a convex domain, the $\bar{\partial}$ -Neumann operator is compact if and only if the boundary contains no q -dimensional analytic varieties. This motivates the following theorem.

Theorem 3. *Let Ω be a bounded convex domain in $\mathbb{C}^n$. Then*

$$(34) \quad A_q = \overline{\bigcup_{\alpha \in \Lambda} f_{\alpha}(\mathbb{D}^q)}.$$

Proof. We first show that if $P \in b\Omega$ is not in A_q , then it is not in the right hand side of (34). Choose $r > 0$ small enough so that $\Omega_1 := \Omega \cap B(P, r)$ is a convex domain whose closure does not intersect A_q . It suffices to establish a compactness estimate on Ω_1 : the result of Fu and Straube mentioned above then implies that the boundary of Ω_1 contains no q -dimensional analytic variety, whence $P \notin \overline{\bigcup_{\alpha} f_{\alpha}(\mathbb{D}^q)}$.

Let $M > 0$. Choose $\varphi_M \in C_0^{\infty}(\mathbb{C}^n)$, $0 \leq \varphi_M \leq 1$, and supported on the set where $-1/M < |z - P|^2 - r^2 < 1/M$. Now let $u \in \text{Dom}(\bar{\partial}) \cap \text{Dom}(\bar{\partial}^*)$ on Ω_1 . Denote by $\lambda_M(z)$ a smooth function that on the support of φ_M agrees with $M(|z - P|^2 - r^2)$ and otherwise is between -1 and 1 . Note that on the support of φ_M , the complex Hessian of λ_M is at least M . To estimate the norm of $\varphi_M u$, we use inequality (2-10) from [5] which says that

$$(35) \quad \sum_K' \sum_{j,k} \int_{\Omega_1} e^{\lambda_M} \frac{\partial^2 \lambda_M}{\partial z_j \partial \bar{z}_k} (\varphi_M u)_{jK} \overline{(\varphi_M u)_{kK}} \leq \|\bar{\partial}(\varphi_M u)\|_{\Omega_1}^2 + \|\bar{\partial}^*(\varphi_M u)\|_{\Omega_1}^2.$$

A comment is in order. (2-10) in [5] is stated for sufficiently smooth domains. We make no smoothness assumptions on Ω other than what is dictated by convexity (Ω is Lipschitz). In addition, Ω_1 has a nonsmooth part in the boundary coming from the intersection of Ω with a small ball. However, the exhaustion procedure

developed in [32] that uses the $\bar{\partial}$ -Neumann operators on a sequence of subdomains allows to forgo any boundary regularity assumptions: (2-10) in [5] holds on any bounded pseudoconvex domain. This is part (ii) of Corollary 1.12 in [34].

For M big enough, we can choose $\chi_M \in C_0^\infty(B(P, r))$, identically equal to one on a neighborhood of the part of the support of $(1 - \varphi_M)$ that lies in $\overline{\Omega_1}$. Note that χ_M (continued by zero outside $B(P, r)$) is a compactness multiplier on Ω (since it vanishes on A_q). Also, the left hand side of (35) dominates qM/e times $\|\varphi_M u\|_{\Omega_1}^2$; the factor q occurs because each term $|u_J|^2$ arises precisely q times as a term $|u_{j,K}|^2$. Therefore, we have for any ε'

$$(36) \quad \begin{aligned} \|u\|_{\Omega_1}^2 &\lesssim \|\varphi_M u\|_{\Omega_1}^2 + \|\chi_M(1 - \varphi_M)u\|_{\Omega}^2 \\ &\lesssim \left(\frac{1}{M} + \varepsilon'\right) \left(\|\bar{\partial}u\|_{\Omega_1}^2 + \|\bar{\partial}^*u\|_{\Omega_1}^2 + \|D\varphi_M u\|_{\Omega}^2\right) + C_{\varepsilon', M} \|(1 - \varphi_M)u\|_{-1, \Omega}^2, \end{aligned}$$

where $D\varphi_M$ denotes a derivative of φ_M . We have used that $(1 - \varphi_M)u \in \text{Dom}(\bar{\partial}) \cap \text{Dom}(\bar{\partial}^*)$ on Ω . To estimate $\|D\varphi_M u\|_{\Omega}^2$, we use again that χ_M is a compactness multiplier on Ω :

$$(37) \quad \begin{aligned} \|D\varphi_M u\|_{\Omega}^2 &= \|\chi_M D\varphi_M u\|_{\Omega}^2 \\ &\leq \varepsilon' C_M \left(\|\bar{\partial}u\|_{\Omega_1}^2 + \|\bar{\partial}^*u\|_{\Omega_1}^2 + \|u\|_{\Omega_1}^2\right) + C_{\varepsilon'} \|D\varphi_M u\|_{-1, \Omega}^2 \\ &\lesssim \varepsilon' C_M \left(\|\bar{\partial}u\|_{\Omega_1}^2 + \|\bar{\partial}^*u\|_{\Omega_1}^2\right) + C_{\varepsilon'} \|D\varphi_M u\|_{-1, \Omega}^2. \end{aligned}$$

Because $(1 - \varphi_M)$ and $D\varphi_M$ are compactly supported in $B(P, r) \cap \overline{\Omega}$, the (-1) -norms on Ω on the right hand side of (36) and (37) are dominated by the corresponding (-1) -norms on Ω_1 and hence by $\|u\|_{-1, \Omega_1}$. Therefore, the desired compactness estimate on Ω_1 results from (36) and (37) upon taking M big enough and then ε' small enough.

For the other direction, assume that $P \notin \overline{\bigcup_{\alpha} f_{\alpha}(\mathbb{D}^q)}$. Choose $r > 0$ small enough so that the closure of $\Omega_1 := B(P, r) \cap \Omega$ is disjoint from $\overline{\bigcup_{\alpha} f_{\alpha}(\mathbb{D}^q)}$. Ω_1 is a convex domain without q -dimensional varieties in the boundary. Again by the Fu-Straube result mentioned above, the $\bar{\partial}$ -Neumann operator on $(0, q)$ -forms on Ω_1 is compact. Therefore, for a smooth function φ supported near P , we have for any $\varepsilon > 0$

$$(38) \quad \begin{aligned} \|\varphi u\|_{\Omega}^2 &= \|\varphi u\|_{\Omega_1}^2 \\ &\leq \varepsilon \left(\|\bar{\partial}(\varphi u)\|_{\Omega_1}^2 + \|\bar{\partial}^*(\varphi u)\|_{\Omega_1}^2\right) + C_{\varepsilon} \|\varphi u\|_{-1, \Omega_1}^2 \\ &\lesssim \varepsilon C_{\varphi} \left(\|\bar{\partial}u\|_{\Omega}^2 + \|\bar{\partial}^*u\|_{\Omega}^2 + \|u\|_{\Omega}^2\right) + C_{\varepsilon, \varphi} \|u\|_{-1, \Omega_1}^2 \\ &\lesssim \varepsilon C_{\varphi} \left(\|\bar{\partial}u\|_{\Omega}^2 + \|\bar{\partial}^*u\|_{\Omega}^2\right) + C_{\varepsilon, \varphi} \|u\|_{-1, \Omega}^2. \end{aligned}$$

In the last inequality, we have used the easily verified inequality $\|u\|_{-1, \Omega_1} \leq \|u\|_{-1, \Omega}$. (38) shows that any such φ is a compactness multiplier on Ω . Choosing a φ that does not vanish at P shows that $P \notin A_q$. This completes the proof of Theorem 3. $\square$

We now turn to complete Hartogs domains in $\mathbb{C}^2$. A complete Hartogs domain Ω in $\mathbb{C}^2$ is defined by $|w| < e^{-\phi(z)}$ for $z \in U$, where U is a domain in $\mathbb{C}$ and $\phi(z)$ is an upper semi-continuous function on U . Ω is pseudoconvex if and only if $\phi(z)$ is subharmonic. If ϕ is at least C^2 , then a computation shows that the weakly pseudoconvex boundary points (z, w) with $w \neq 0$ are those where $\Delta\phi(z) = 0$ (see also [29], p. 100).

On a smooth bounded pseudoconvex (not necessarily complete) Hartogs domain compactness of the $\bar{\partial}$ -Neumann operator is equivalent to Catlin's property (P) ([10]). Additionally, if the domain is also complete and the boundary points with $w = 0$ are strictly pseudoconvex, then both of the above conditions (compactness and property (P)) are equivalent to the following: the projection of the weakly pseudoconvex boundary points into the z -plane has empty fine interior ([31], p.310, [34], Lemma 3.19). Recall that the fine topology is the smallest topology that makes all subharmonic functions continuous; see, e.g. [18, 1] for properties of this topology. It is strictly larger than the Euclidean topology, and there exist compact sets with empty Euclidean interior, but nonempty fine interior, see [1] example 7.9.3. Of importance here will be the following fact, which explains why the fine topology plays a role in our context: a compact subset of the plane satisfies property (P) if and only if it has empty fine interior ([31], Proposition 1.11, [34], Proposition 3.17). We will use the notation $Int_f(K)$ for the fine interior points of the set K , and $\bar{B}^E$ for the Euclidean closure of a set B .

Denote by π the projection $\pi : \mathbb{C}^2 \rightarrow \mathbb{C}$. $(z, w) \rightarrow z$. Let K be the projection of the weakly pseudoconvex boundary points. When the boundary points with $w = 0$ are strictly pseudoconvex, $K = \{z \in U \mid \Delta\phi(z) = 0\}$, and K is a compact subset of the base domain U . On such a domain we have the following characterization of $A := A_1$.

Theorem 4. *Let Ω be a smooth bounded complete pseudoconvex Hartogs domain in $\mathbb{C}^2$. Assume that the boundary points of the form $(z, 0)$ are strictly pseudoconvex. Then*

$$A = \overline{\pi^{-1}(Int_f(K))}^E.$$

Proof. The structure of the proof is the same as in the proof of Theorem 3, but the details change. We first show that if $P \in b\Omega \setminus \overline{\pi^{-1}(Int_f(K))}^E$, then $P \in b\Omega \setminus A$. Choose $r > 0$ small enough so that $\pi(B(P, r))$ is disjoint from $\overline{Int_f(K)}^E$ and set $\Omega_1 := B(P, r) \cap \Omega$. We will show that $b\Omega_1$ satisfies property (P_1) , so that the $\bar{\partial}$ -Neumann operator on Ω_1 is compact. The rest of the argument then follows that in the second part of the proof of Theorem 3.

We write $b\Omega_1$ as a countable union of compact sets, all of which satisfy property (P_1) ; then so does $b\Omega_1$ ([31], Proposition 1.9, [34], Corollary 3.14). The first set is $bB(P, r) \cap \bar{\Omega}$. The second set consists of the set W of weakly pseudoconvex boundary points of Ω that are contained in $\bar{B}(P, r)$. Note that $\pi(W)$ has empty fine interior, hence satisfies property (P_1) in the plane. Therefore, W satisfies property (P_1) , by [31], Proposition 1.10 (the image $\pi(W)$ does, and the fiber over each point is a circle

and so also satisfies (P_1) . Finally, write the set of strictly pseudoconvex boundary points of Ω as a countable union of compact sets K_n ; they then satisfy property (P_1) . Thus so do the compacts $K_n \cap \overline{B(P, r)}$. The union of these sets together with W and the first set above equals $b\Omega_1$, and we are done.

Now let $P \in b\Omega \setminus A$. We will construct a Hartogs domain Ω_1 that shares a (rotationally invariant) piece of boundary with Ω that contains P and does not intersect A . In addition, Ω_1 will be strictly pseudoconvex off of that shared piece. Observe that A is invariant under rotations in w : pullbacks commute with $\bar{\partial}$ and, because the rotations induce isometries on $L^2_{(0,1)}(\Omega)$, also with $\bar{\partial}^*$ (this observation was exploited in [4]). So f is a compactness multiplier if and only if $f_\theta(z, w) := f(z, e^{i\theta}w)$ is. Therefore, $P \in b\Omega \setminus A$ implies $P_1 := \pi(P) \notin \pi(A)$. Choose radii $0 < r_1 < r_2 < r_3$ such that $\overline{D(P_1, r_3)} \cap \pi(A) = \emptyset$. Ω_1 is going to be over the base $D(P_1, r_3)$. Then choose $\varphi(z) \in C_0^\infty(D(P_1, r_3))$ such that $\varphi(z) \equiv 1$ on $\overline{D(P_1, r_2)}$. Now set $\psi(z) := \phi(z)\varphi(z) + h(z)$ on $D(P_1, r_3)$, where $h(z)$ is a smooth radially symmetric subharmonic function on $D(P_1, r_3)$, $h(z) = 0$ on $\overline{D(P_1, r_1)}$, and equal to $-\frac{1}{2}\log(r_3^2 - |z - P_1|^2)$ when $|z - P_1|$ is close to r_3 . Such a function can be chosen to be increasing and concave up, and to have its second (radial) derivative as big as we wish on a given compact subset of $D(P_1, r_3) \setminus \overline{D(P_1, r_1)}$, in particular on $\{\Delta(\phi\varphi) \leq 0\} \cap \text{supp}(\varphi) \cap \{|z| \geq r_2\}$. That means that the Laplacian of h on this set can be made as big as we wish. Making this Laplacian big enough ensures that $\psi(z)$ is subharmonic, and

$$\Omega_1 := \{(z, w) \in \mathbf{C}^2 \mid z \in D(P_1, r_3), |w| < e^{-\psi(z)}\}$$

is a smooth bounded pseudoconvex complete Hartogs domain which near the boundary of its base looks like a ball. This construction comes from [17], proof of Theorem 4.2; see also [34], proof of Theorem 3.25.

We claim that the portion of the boundary of Ω_1 that sits over $\{r_1 \leq |z - P_1| \leq r_3\}$ satisfies property (P_1) . Parts over compact subsets of $\{r_1 < |z - P_1| < r_3\}$ are compact subsets of the strictly pseudoconvex part of the boundary and so satisfy property (P_1) . The part corresponding to $\pi^{-1}(\{|z| = r_1\})$ satisfies (P_1) for the same reason that the set W in the first part of the proof did (i.e. by [31], Proposition 1.10). The circle $\{|z| = r_3\}$ also satisfies (P_1) . Thus the portion of the boundary we are interested in is the countable union of compact sets that satisfy property (P_1) , and the claim is established (again as in the first part of the proof).

Using the claim from the previous paragraph together with the fact that for each boundary point that is common to Ω_1 and Ω (these are the boundary points of Ω_1 over the set $\{|z| \leq r_1\}$) there exists a compactness multiplier on Ω that does not vanish at the point, one can follow the first part of the proof of Theorem 3 to show that the $\bar{\partial}$ -Neumann problem on Ω_1 satisfies a compactness estimate. By the result of Christ and Fu [10], the boundary of Ω_1 satisfies property (P_1) , and by our discussion above, the projection of its weakly pseudoconvex boundary points onto the z -plane has empty fine interior. Therefore (because Ω and Ω_1 share an open piece

of boundary near P), $P_1 \notin \overline{\text{int}_f(K)}^E$, and consequently $P \in b\Omega \setminus \overline{\pi^{-1}(\text{Int}_f(K))}^E$. This completes the proof of Theorem 4. $\square$

REFERENCES

- [1] D. H. Armitage and S. J. Gardiner, *Classical Potential Theory*, Springer Monographs in Mathematics, Springer-Verlag London Ltd., London, 2001.
- [2] D. E. Barrett, *Behavior of the Bergman projection on the Diederich-Fornæss worm*, Acta Math. **168** (1992), no. 1-2, 1–10.
- [3] H. P. Boas, *Small sets of infinite type are benign for the $\bar{\partial}$ -Neumann problem*, Proc. Amer. Math. Soc. **103**, no.2 (1988), 569–578.
- [4] H. P. Boas, S. Ch. Chen, and E. J. Straube, *Exact regularity of Bergman and Szegő projections on domains with partially transverse symmetries*, Manuscripta Math. **62** (1988), 467–475.
- [5] H. P. Boas and E. J. Straube, *Global regularity of the $\bar{\partial}$ -Neumann problem: a survey of the L^2 -Sobolev theory*, Several Complex Variables (Berkeley, CA, 1995–1996), Math. Sci. Res. Inst. Publ., vol. 37, Cambridge Univ. Press, Cambridge, 1999, pp. 79–111.
- [6] D. W. Catlin, *Global regularity of the $\bar{\partial}$ -Neumann problem*, Complex Analysis of Several Variables, Proc. Sympos. Pure Math., vol. 41, Amer. Math. Soc., Providence, RI, 1984, pp. 39–49.
- [7] ———, *Subelliptic estimates for the $\bar{\partial}$ -Neumann problem on pseudoconvex domains*, Ann. Math. (2) **126** (1987), 131–191.
- [8] M. Çelik, *Contributions to the compactness theory of the $\bar{\partial}$ -Neumann operator*, Ph. D. dissertation, Texas A&M University, May 2008.
- [9] S. C. Chen and M. C. Shaw, *Partial Differential Equations in Several Complex Variables*, AMS/IP Studies in Advanced Mathematics, vol. 19, American Mathematical Society, Providence, RI, 2001.
- [10] M. Christ and S. Fu, *Compactness in the $\bar{\partial}$ -Neumann problem, magnetic Schrödinger operators, and the Aharonov-Bohm effect*, Adv. Math. **197** (2005), no. 1, 1–40.
- [11] J. P. D’Angelo, *Several Complex Variables and the Geometry of Real Hypersurfaces*, Studies in Advanced Mathematics, CRC Press, 1993.
- [12] ———, *Inequalities from Complex Analysis*, Carus Mathematical Monographs, 28. Mathematical Association of America, Washington, DC, 2002.
- [13] J. P. D’Angelo and J. J. Kohn, *Subelliptic estimates and finite type*, Several Complex Variables, Math. Sci. Res. Inst. Publ., vol. 37, Cambridge Univ. Press, Cambridge, 1999, pp. 199–232.
- [14] E. B. Davies, *Spectral Theory and Differential Operators*, Cambridge Studies in Advanced Mathematics, vol. 42, Cambridge University Press, 1995.
- [15] G. B. Folland and J. J. Kohn, *The Neumann Problem for the Cauchy-Riemann Complex*, Princeton University Press, Princeton, N.J., 1972, Annals of Mathematics Studies, No. 75.
- [16] S. Fu and E. J. Straube, *Compactness of the $\bar{\partial}$ -Neumann problem on convex domains*, J. Funct. Anal. **159** (1998), no. 2, 629–641.
- [17] ———, *Compactness in the $\bar{\partial}$ -Neumann problem*, Complex Analysis and Geometry, Ohio State Univ. Math. Res. Inst. Publ., vol. 9, de Gruyter, Berlin, 2001, pp. 141–160.
- [18] L. L. Helms, *Introduction to Potential Theory*, Pure and Applied Mathematics, Vol. XXII, Wiley-Interscience A Division of John Wiley & Sons, New York-London-Sydney, 1969.
- [19] L. Hörmander, *L^2 estimates and existence theorems for the $\bar{\partial}$ operator*, Acta Math. **113** (1965), 89–152.
- [20] J. J. Kohn, *Global regularity for $\bar{\partial}$ on weakly pseudo-convex manifolds*, Trans. Amer. Math. Soc. **181** (1973), 273–292.
- [21] ———, *Subellipticity of the $\bar{\partial}$ -Neumann problem on pseudo-convex domains: sufficient conditions*, Acta Math. **142** (1979), no. 1-2, 79–122.
- [22] J. J. Kohn and L. Nirenberg, *Non-coercive boundary value problems*, Comm. Pure Appl. Math. **18** (1965), 443–492.

- [23] S. Lang, *Real and Functional Analysis*, third ed., Graduate Texts in Mathematics, vol. 142, Springer-Verlag, New York, 1993.
- [24] J. L. Lions and E. Magenes, *Non-Homogeneous Boundary Value Problems and Applications. Vol. I*, Springer-Verlag, New York, 1972, Translated from the French by P. Kenneth, Die Grundlehren der mathematischen Wissenschaften, Band 181.
- [25] P. Matheos, *A Hartogs domain with no analytic disc in the boundary for which the $\bar{\partial}$ -Neumann problem is not compact*, Ph. D. dissertation, University of California, Los Angeles, 1997.
- [26] J. D. McNeal, *A sufficient condition for compactness of the $\bar{\partial}$ -Neumann operator*, J. Funct. Anal. **195** (2002), no. 1, 190–205.
- [27] A. Nicoara, *Equivalence of types and Catlin boundary systems*, preprint.
- [28] R. M. Range, *The $\bar{\partial}$ -Neumann operator on the unit ball in $\mathbb{C}^n$* , Math. Ann. **266** (1984), 449–456.
- [29] ———, *Holomorphic Functions and Integral Representations in Several Complex Variables*, Graduate Texts in Mathematics, vol. 108, Springer-Verlag, New York, 1986.
- [30] M. Reed and B. Simon, *Methods of Modern Mathematical Physics. I*, second ed., Academic Press Inc., New York, 1980, Functional analysis.
- [31] N. Sibony, *Une classe de domaines pseudoconvexes*, Duke Math. J. **55** (1987), no. 2, 299–319.
- [32] E. J. Straube, *Plurisubharmonic functions and subellipticity of the $\bar{\partial}$ -Neumann problem on non-smooth domains*, Math. Res. Lett. **4** (1997), no. 4, 459–467.
- [33] ———, *Aspects of the L^2 -Sobolev theory of the $\bar{\partial}$ -Neumann problem*, International Congress of Mathematicians. Vol. II, Eur. Math. Soc., Zürich, 2006, pp. 1453–1478.
- [34] ———, *Lectures on the L^2 -Sobolev Theory of the $\bar{\partial}$ -Neumann Problem*, Monograph in preparation, 2008.
- [35] W. J. Sweeney, *Coerciveness in the Neumann problem*, J. Differential Geometry **6** (1971/72), 375–393.

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843
 E-mail address: celik@math.tamu.edu, straube@math.tamu.edu